arXiv:hep-th/0506050v2 30 Jun 2005

UTAP-528, RESCEU-9/05

Warped Flux Compactification and Brane Gravity

Shinji Mukohyama[a,b], Yuuiti Sendouda[a],
Hiroyuki Yoshiguchi[a], Shunichiro Kinoshita[a]

[a]*Department of Physics*
The University of Tokyo, Tokyo 113-0033, Japan

[b]*Research Center for the Early Universe*
The University of Tokyo, Tokyo 113-0033, Japan

Abstract

We find a simple exact solution of 6-dimensional braneworld which captures some essential features of warped flux compactification, including a warped geometry, compactification, a magnetic flux, and one or two 3-brane(s). In this setup we analyze how the Hubble expansion rate on each brane changes when the brane tension changes. It is shown that effective Newton's constant resulting from this analysis agrees with that inferred by simply integrating extra dimensions out. Based on the result, a general formula for effective Newton's constant is conjectured and its application to cosmology with type IIB warped string compactification is discussed.

1 Introduction

Warped flux compactification is one of essential parts of the construction of de Sitter vacua in string theory by Kachru, Kallosh, Linde and Trivedi [1] [1]. A warped throat region [11] is smoothly attached to a Calabi-Yau manifold and all moduli except for the volume modulus are stabilized by fluxes [12]. The volume modulus is thought to be stabilized by non-perturbative effects such as D-instantons [13, 1]. Another important ingredient of the construction is branes. Anti-D-branes are located at the bottom of the throat region to uplift the stable AdS vacua to meta-stable de Sitter vacua. This setup provides a number of possible applications to cosmology [14, 15, 16, 17, 18].

So far, there is no known explicit form of the global geometry including the throat region, compactification, fluxes and the moduli stabilization. The lack of known explicit form of the global geometry makes it difficult to analyze brane gravity in the context of warped flux compactification in details from higher dimensional point of view. Of course, even without the explicit global geometry, one could invoke that the 4-dimensional Einstein gravity should be recovered as a low energy effective theory since all moduli are thought to be stabilized and made massive so that they can be integrated out. Nonetheless, it is perhaps fair to say that we have not yet had a complete understanding of how 4-dimensional Einstein gravity is recovered in the warped flux compactification.

For example, suppose that a D-brane and an anti-D-brane annihilate somewhere in the warped throat region but that they are not coincident with a brane on which we are living. Following the idea of [19, 20], it is suggested that this process may drive an inflation in our 4-dimensional universe [14] and could leave cosmic superstrings as relics [17]. In this picture the inflaton and cosmic superstrings are not on our brane but living somewhere in the extra dimensions. It does not seem completely clear how they affect gravity on our brane. Indeed, a lot of questions would arise. “Does the inflaton living outside our world really inflate our brane?” “Is there a deficit angle on our brane due to the cosmic superstring wandering somewhere in the extra dimensions?” “If there is a deficit angle, then where on our brane and how much?” Partial answers to these questions would be obtained if we would completely integrate out extra degrees of freedom due to the extra dimensions, provided that all moduli are stabilized. On the other hand, in principle it should also be possible to answer these questions by analyzing the higher dimensional theory directly. Evidently, it is important to take both approaches complementarily and compare them towards our good understanding of brane gravity in the warped flux compactification.

---

[1] See [2, 3] for followup proposals. See also [4, 5], [6, 7, 8, 9] and [10] for other proposals of de Sitter, transiently accelerating and inflationary universes.

Now let us heuristically remind ourselves how 4-dimensional Einstein gravity is recovered in the Kaluza-Klein compactification and the Randall-Sundrum type braneworlds.

In Kaluza-Klein compactification, zero modes and Kaluza-Klein (KK) modes are decoupled in the linearized level because of the momentum conservation along extra dimensions. This simple fact makes the recovery of the 4-dimensional theory manifest since the standard model fields are supposed to consist of zero modes. Indeed, as far as moduli associated with compactification are made massive by a stabilization mechanism, they do not appear in the low energy physics and the Einstein theory is recovered as a low energy effective theory. This consideration also applies to cosmology as far as the energy scale is sufficiently lower than the compactification scale.

In the second Randall-Sundrum (RS2) braneworld with infinite extra dimension [21], the recovery of the 4-dimensional Einstein theory is due to localization of zero mode [21, 22, 23, 24]. Matter on our brane possibly excites not only zero modes but also KK modes since the momentum conservation does not prohibit couplings between the singular brane source and the KK modes [2]. However, the warped geometry localizes the zero mode to the vicinity of the brane and the coupling of the brane source to the zero mode is much stronger than that to KK modes. In this way, gravity on our brane at low energy is almost locally determined by the zero mode localized near the brane and the 4-dimensional Einstein theory is recovered at low energy. (This point of "locally localized gravity" has been made explicit in ref. [25].) For FRW cosmology on the brane, there is no well-defined distinction between zero modes and KK modes because of the lack of enough symmetry. Nonetheless, the evolution of the brane is still determined locally since the unbroken symmetry, i.e. the homogeneity and the isotropy parallel to the brane, prevents waves from being generated and propagating in the bulk. In this way, gravity is still localized and the standard cosmological equation is recovered at low energy [26, 27, 28, 29, 30, 31, 34, 32, 33].

On the other hand, in the warped compactification the recovery of 4-dimensional theory seems more indirect and subtle. Unlike KK compactification but as in RS2 braneworld, matter source on the brane can excite not only zero modes but also KK modes. However, unlike the RS2 braneworld, gravity is not localized near the brane since the warp factor on the brane in the throat region is not larger but smaller than that in the bulk nearby. Hence, the evolution of matter on the brane changes the bulk geometry not only near the brane but possibly everywhere in the whole extra dimensions. Nonetheless, if all moduli are stabilized, the bulk geometry should quickly settle to a configuration which is determined by the boundary condition, i.e. the brane source(s), values of conserved quantities and the regularity of the other region of the extra dimensions. As a consequence of the change of the bulk geometry, the induced geometry on the brane responses to the evolution of the matter source on the brane. It is quite possible but must be checked that the 4-dimensional Einstein theory is recovered as a rather indirect and subtle relation between the matter source on the brane and the response of the induced geometry. In this paper we shall support this picture in a simplified situation.

---

[2] Note that a delta function includes all momenta when it is Fourier transformed.

This picture is somehow similar to that in the first Randall-Sundrum (RS1) scenario [35] with radion stabilization [36]. In the original RS1 brane model (without radion stabilization) the 4-dimensional theory is not Einstein but Brans-Dicke theory [22, 23] because of the existence of a massless modulus called radion. With the radion stabilization, there is no extra massless degree which could appear in the 4-dimensional effective theory and the 4-dimensional Einstein theory is recovered at energies sufficiently below the stabilization scale. The recovery of the 4-dimensional Einstein theory is understood both from the low energy effective theory point of view and as a consequence of dynamics of bulk fields [37, 38, 39, 40, 41, 42, 43].

In the case of KK compactification and RS braneworlds, the recovery of the Einstein theory has been explicitly investigated from higher dimensional point of view. One of the reasons why this was possible is that there are explicit background solutions (or at least explicit equations defining background solutions) around which we can analyze perturbations. On the other hand, in the warped flux compactification, the absence of an explicit global solution makes it less tractable to see the recovery of the 4-dimensional Einstein gravity from higher dimensional point of view as explicitly as in the KK compactification and the RS braneworlds.

Therefore, we would like to consider a simplified situation in which we can see the recovery of the 4-dimensional Einstein theory in the warped flux compactification. The purpose of this paper is, as a first step, to consider a toy model which captures some essential features of the warped flux compactification and to see explicitly how the induced geometry on a brane responses to brane tension as a consequence of changes in the bulk geometry. In particular, we shall see that the relation between the change of brane tension and the response of the induced geometry is identical to that inferred from the 4-dimensional Einstein theory.

The rest of this paper is organized as follows. In Sec. 2 we describes a simple model of 6-dimensional warped flux compactification with one or two 3-brane(s). In Sec. 3 we argue that the 4-dimensional Friedmann equation should be recovered on each brane at low energy and confirms the validity of a formula of effective Newton's constant in a simplified situation. Sec. 4 is devoted to a summary of this paper and discussion. In particular, based on the result of this paper, a general formula for effective Newton's constant is conjectured and its application to cosmology with type IIB warped string compactification is discussed. In Appendices A.1 and A.2 we show that two seemingly singular limits of the solutions considered in this paper are actually regular. In Appendix A.3 a higher-order correction to the effective Friedmann equation on a brane is estimated and it is shown that higher order corrections can be ignored when the Hubble expansion rate on the brane is sufficiently lower than the bulk curvature scale.

2 Model description

In this paper we would like to consider a minimal setup which captures essential features of the warped flux compactification and which is simple enough to analyze brane gravity from higher dimensional viewpoint. The setup must include at least a warped geometry, magnetic flux of an antisymmetric field along the extra dimensions and a brane. Since the simplest antisymmetric field is a $U(1)$ gauge field and the corresponding magnetic flux has two spatial indices, we need to consider at least 2 extra dimensions and, thus, at least 6-dimensional spacetime. In this section we shall describe a model of 6-dimensional braneworld with warped flux compactification. In the next section we shall analyze this model in a situation where the tension of a 3-brane changes by a phase transition on the brane.

For the reason explained above, in this paper we consider a 6-dimensional braneworld scenario with a $U(1)$ gauge field and a cosmological constant in the bulk. The bulk action is

$$I_6 = \frac{M_6^4}{2} \int d^6x\sqrt{-g}\left(R - 2\Lambda_6 - \frac{1}{2}F^{MN}F_{MN}\right), \tag{2.1}$$

where M_6 is the 6-dimensional reduced Planck mass, Λ_6 is the bulk cosmological constant, and $F_{MN} = \partial_M A_N - \partial_N A_M$ is the field strength associated with the $U(1)$ gauge field A_M.

2.1 Bulk solution

We assume that the 6-dimensional bulk geometry has the 4-dimensional de Sitter symmetry and an additional axisymmetry. The former symmetry is to make the induced geometry on our brane a de Sitter spacetime, and the latter is imposed as a symmetry in the extra dimensions for simplicity. The general metric with these symmetries is locally written as

$$ds_6^2 = A(w)^2 ds_4^2 + dw^2 + R(w)^2 d\phi^2, \tag{2.2}$$

where w is the proper distance along geodesics orthogonal to the brane world-volume and to the orbit of the axisymmetry, $A(w)$ and $R(w)$ are functions of w only, and ds_4^2 is the line element of the 4-dimensional de Sitter spacetime with the unit curvature radius. For example, in the global chart ds_4^2 is written as

$$ds_4^2 = -dt^2 + \cosh^2 t d\Omega_3^2, \tag{2.3}$$

where $d\Omega_3^2$ is the line element of the unit 3-sphere.

If $A(w)$ is constant then the solution to the Einstein equation is of the ADD type [44]. This solution was already investigated in [45, 46, 47]. In Appendix A.1, we shall see that this solution is a particular limit of the solutions below.

Our interest is in the case of non-vanishing $\partial_w A$ since this corresponds to a warped geometry [3]. (As already stated, we would like to consider a minimal setup which includes at least a warped geometry, a flux and a brane.) In this case we can introduce a new coordinate r by $r = A(w)$ at least locally. With the new coordinate the line element is

$$ds_6^2 = r^2 ds_4^2 + g(r)dr^2 + f(r)d\phi^2, \tag{2.4}$$

where $f(r)$ and $g(r)$ are functions of r only. It is of course possible and indeed straightforward to analyze the Einstein equation and find solutions with this ansatz. It is also possible to take double Wick rotation, find a family of solutions and then take inverse double Wick rotation. By the double Wick rotation

$$t \to i\left(\frac{\pi}{2} - \theta\right), \quad \phi \to iT, \tag{2.5}$$

the metric ansatz (2.4) is transformed to

$$d\tilde{s}_6^2 = -\tilde{f}(r)dT^2 + \tilde{g}(r)dr^2 + r^2 d\Omega_4^2, \tag{2.6}$$

where

$$d\Omega_4^2 = d\theta^2 + \sin^2\theta d\Omega_3^2 \tag{2.7}$$

is the line element of the unit 4-sphere. The metric (2.6) is nothing but a general ansatz for a spherically symmetric, static metric. There is a well-known family of solutions: RN-de Sitter (for $\Lambda_6 > 0$), RN (for $\Lambda_6 = 0$) and RN-AdS (for $\Lambda_6 < 0$) spacetime. The RN-de Sitter, RN or RN-AdS solution is

---

[3] This setup was considered in [48, 49] but, as far as we know, the family of explicit exact solutions of warped flux compactification presented below had not been found in the literature.

$$\begin{aligned} \tilde{f}(r) &= \frac{1}{\tilde{g}(r)} = 1 - \frac{\Lambda_6}{10}r^2 - \frac{\mu}{r^3} + \frac{e^2}{12r^6}, \\ A_M dx^M &= \frac{e}{3r^3}dT, \end{aligned} \tag{2.8}$$

where μ is a constant of integration corresponding to the mass parameter and e is the electric charge. Going back to the original ansatz by the inverse double Wick rotation

$$\frac{\pi}{2} - \theta \to -it, \quad T \to -i\phi, \tag{2.9}$$

we obtain the family of solutions

$$\begin{aligned} f(r) &= \frac{1}{g(r)} = 1 - \frac{\Lambda_6}{10}r^2 - \frac{\mu}{r^3} - \frac{b^2}{12r^6}, \\ A_M dx^M &= \frac{b}{3r^3}d\phi, \end{aligned} \tag{2.10}$$

where we have introduced the magnetic charge b by $e \to ib$ so that $A_M dx^M$ remains real.

2.2 Brane sources

In the following we shall consider one or two 3-brane sources. For this purpose we shall use the well-known formula

$$\delta_\pm = \frac{\sigma_\pm}{M_6^4}, \tag{2.11}$$

where $\delta_\pm$ is the deficit angle due to the tension $\sigma_\pm$ of the branes. We, however, have to keep in mind that there is no simple general prescription, analogous to Israel's junction condition [50], for obtaining physical characteristics of an arbitrary distributional source with more than one codimensions [51, 52]. The difficulty is essentially due to the fact that Einstein equation is non-linear. The formula (2.11) is valid under the axisymmetry if radial stress is much smaller than energy density [53]. Namely, with these conditions, $\sigma_\pm$ in (2.11) can be effectively considered as inertial mass per brane's volume. On the other hand, if radial stress is not negligible then the formula (2.11) should be considered as the definition of $\sigma_\pm$, which is in general different from inertial mass per brane's volume.

Hereafter we assume that the function $f(r)$ given by (2.10) has two positive roots $r = r_\pm$ ($0 < r_- < r_+$) and is positive between them ($r_- < r < r_+$). This requires that the 6-dimensional cosmological constant Λ_6 be positive. Thus, hereafter we assume that $\Lambda_6 > 0$. Since f vanishes at $r = r_\pm$, the equation $r = r_\pm$ defines surfaces of codimension 2.

Near $r = r_\pm$ in the bulk ($r_- < r < r_+$),

$$\frac{dr^2}{f(r)} + f(r)d\phi^2 \simeq d\rho_\pm^2 + \kappa_\pm^2 \rho_\pm^2 d\phi^2, \tag{2.12}$$

where

$$\rho_\pm \equiv \sqrt{\frac{\mp 2(r - r_\pm)}{\kappa_\pm}}, \quad \kappa_\pm \equiv \mp\frac{1}{2}f'(r_\pm)\ (>0). \tag{2.13}$$

Hence, with the deficit angles $\delta_\pm$ at $r = r_\pm$, respectively, ϕ is identified as

$$\kappa_\pm \phi \sim \kappa_\pm \phi + (2\pi - \delta_\pm). \tag{2.14}$$

The identification at $r = r_+$ is consistent with that at $r = r_-$ if and only if

$$\frac{2\pi - \delta_+}{2\pi - \delta_-} = \frac{\kappa_+}{\kappa_-}. \tag{2.15}$$

This can be considered as a boundary condition since the l.h.s. is specified by the brane sources and the r.h.s. can be written in terms of the bulk parameters μ and b.

In this way, we can put 3-branes at $r = r_\pm$ and consider the region $r_- < r < r_+$ as the bulk spacetime. It is also possible to consider a solution with only one 3-brane at either $r = r_+$ or $r = r_-$ by setting $\delta_- = 0$ or $\delta_+ = 0$. The Hubble parameter $H_\pm$ on the brane at $r = r_\pm$ is given by

$$H_\pm = \frac{1}{r_\pm}. \tag{2.16}$$

3 Recovery of the Friedmann equation

Let us consider a $(4+n)$-dimensional, general warped compactification

$$ds_{4+n}^2 = r^2 g_{\mu\nu}^{(4)} dx^\mu dx^\nu + \gamma_{ij} dy^i dy^j, \tag{3.1}$$

where the 4-dimensional metric $g_{\mu\nu}^{(4)}$ depends on the 4-dimensional coordinates x^μ only, and the n-dimensional metric γ_{ij} and the warp factor r depend only on the coordinates y^i of the compact extra dimensions. With this warped metric, the $(4+n)$-dimensional Einstein-Hilbert action includes the 4-dimensional Einstein term:

$$(M_{4+n})^{2+n} \int d^4x d^ny \sqrt{-g^{(4+n)}} R^{(4+n)} = (M_{4+n})^{2+n} \int d^ny \sqrt{\gamma}\, r^2 \times \int d^4x \sqrt{-g^{(4)}} R^{(4)} + \cdots, \tag{3.2}$$

where dots represent terms including derivatives of r, the curvature of γ_{ij} and so on. Therefore, if all moduli associated with the extra dimensions are stabilized and if their masses are large enough then we expect that the 4-dimensional Einstein theory is recovered at low energy and that Newton's constant G_N on a brane at $y^i = y^i_0$ should be given by

$$\frac{1}{8\pi G_N} = (M_{4+n})^{2+n} \int d^n y \sqrt{\gamma} \left[\frac{r(y)}{r(y_0)} \right]^2 , \tag{3.3}$$

where the normalization factor $[r(y_0)]^{-2}$ has been included in order to take into account the fact that the induced metric on the brane is not $g^{(4)}_{\mu\nu} dx^\mu dx^\nu|_{y=y_0}$ but $r^2 g^{(4)}_{\mu\nu} dx^\mu dx^\nu|_{y=y_0}$.

This expectation is known to be correct for codimension 1 ($n = 1$) braneworlds with radion stabilization [37, 38, 39, 40, 41, 42, 43] [4].

We expect that the formula (3.3) should be correct also for codimension 2 or higher ($n \geq 2$) braneworlds if all moduli are stabilized. In this paper, for simplicity, we consider the 6-dimensional braneworld described in the previous section and a situation where the tension of one of the 3-branes at $r = r_\pm$ changes by a phase transition on the brane. We suppose that the tension is almost constant in deep inside the old and new phases. With this setup, it is expected that the 4-dimensional geometries on the brane deep inside the two phases are approximated by de Sitter spacetimes with different Hubble expansion rates. What we should see is the relation between the difference of tension in the two phases and the corresponding change of the Hubble expansion rate. Indeed, we shall see below that at low energy, the relation is identical to that inferred from the standard Friedmann equation, provided that 4-dimensional Newton's constant is given by the formula (3.3). We shall also see what "low energy" exactly means.

By the phase transition, the bulk geometry should change since the boundary condition set by the brane tension changes. In particular, the bulk parameters μ and b after the phase transition will in general be different from what they were before. Actually, by the following reason, the values of μ and b after the phase transition should be uniquely determined by the brane tension. Indeed, the boundary condition (2.15) and the conservation of magnetic flux provide two independent conditions on the two independent parameters μ and b and, thus, uniquely fix the parameters at least for a sufficiently small change.

When one of the brane tensions changes, the deficit angle at the position of the brane changes, according to the formula (2.11). This in general induces the change of the area of the extra dimensions, in particular the interval of ϕ. Since the magnetic flux is nothing but the integral of the magnetic field over the extra dimensions, the change of the area and the flux conservation imply that the amplitude b of magnetic field should change. At the same time, the parameters μ and b must satisfy the boundary condition (2.15) with the new deficit angle corresponding to the tension after the phase transition. Therefore, not only b but also μ should change in general. In this way, the values of b and μ after the phase transition are uniquely determined by the flux conservation and the boundary condition.

---

[4] With the Z_2 symmetry, the integration over the bulk must be multiplied by 2 to take into account the fact that there are two copies of the same bulk geometry.

Accordingly, positions $r_\pm$ of branes are also determined uniquely since they are defined as roots of the function $f(r)$ in (2.10). In other words, the Hubble expansion rate $H_\pm = 1/r_\pm$ changes after the phase transition and there is a unique relation between the change of the brane tension and the change of the Hubble expansion rate. Since all relevant equations are invariant under the reflection $H_\pm \to -H_\pm$, the resulting physical relation must also be invariant under this reflection. Therefore, the relation must be even in $H_\pm$ and its Taylor expansion w.r.t. $\Delta\sigma_\pm$ (or equivalently, w.r.t. $\Delta H_\pm^2$) should start as

$$\Delta H_\pm^2 \propto \Delta\sigma_\pm + O((\Delta\sigma_\pm)^2), \tag{3.4}$$

where ΔX represents a small difference of the quantity X before and after the phase transition. Finally, from this relation we obtain

$$H_\pm^2 \propto (\sigma_\pm - \sigma_\pm^{(0)}) + O((\sigma_\pm - \sigma_\pm^{(0)})^2) \tag{3.5}$$

for small $H_\pm^2$, where $\sigma_\pm^{(0)}$ is the value of $\sigma_\pm$ corresponding to $H_\pm^2 = 0$. In this argument, it has been implicitly assumed that the limit $H_\pm^2 \to 0$ is regular. In Appendix A.2 we explicitly confirm that this limit is indeed regular.

The relation (3.5) is nothing but the standard Friedmann equation if the proportionality coefficient is $8\pi G_N/3$, where G_N is Newton's constant. In the following, we shall see both numerically and analytically that this is indeed the case, where Newton's constant is given by the formula (3.3). To be more specific, we shall show that Newton's constant $G_{N\pm}$ on the brane at $r = r_\pm$, respectively, is given by

$$\frac{1}{8\pi G_{N\pm}} = M_6^4 \int_{r_-}^{r_+} \left(\frac{r^2}{r_\pm^2}\right) dr \int_0^{\Delta\phi} d\phi = M_6^4 L^2 \Sigma \cdot \left(\frac{r_-}{r_+}\right)^{\pm 1}, \tag{3.6}$$

where $L \equiv \sqrt{10/\Lambda_6}$,

$$\Sigma \equiv \frac{1}{L^2} \int_{r_-}^{r_+} \frac{r^2}{r_+ r_-} dr \int_0^{\Delta\phi} d\phi = \frac{\Delta\phi}{3L^2 r_+ r_-}(r_+^3 - r_-^3) \tag{3.7}$$

is the warped volume of the extra dimensions (in the unit of L^2), and

$$\Delta\phi = \frac{2\pi - \delta_+}{\kappa_+} = \frac{2\pi - \delta_-}{\kappa_-} \tag{3.8}$$

is the period of the angular coordinate ϕ.

For later convenience, here we define the magnetic flux Φ in the unit of L as

$$\Phi = \frac{1}{L}\int_{r_-}^{r_+} dr \int_0^{\Delta\phi} d\phi F_{r\phi} = -\frac{b\Delta\phi}{3L}\left(\frac{1}{r_-^3} - \frac{1}{r_+^3}\right). \tag{3.9}$$

All relevant equations including this are invariant under the reflection $(b, \Phi) \to (-b, -\Phi)$. Thus, we do not need to keep track of the sign of Φ as far as its sign relative to b is correct. As already stated, the magnetic flux is conserved and, thus, must be fixed during the phase transition.

3.1 Re-parameterization

As already explained, we expect that the Friedmann equation should be recovered at low energy as a consequence of the response of the bulk geometry to the evolution of the brane source. In this respect it is not physically relevant to keep track of the change of the bulk parameters μ and b. What is physically important is, instead, the brane tensions (or equivalently the deficit angles) and the Hubble expansion rates on the branes (or equivalently the positions $r_\pm$ of the branes).

Hence, it is useful to express the metric in terms of $h \equiv L\sqrt{H_+H_-} = L/\sqrt{r_+r_-}$ and $\alpha \equiv H_+/H_- = r_-/r_+$ instead of the original parameters μ and b. By the definition of $r_\pm$, the new parameter α satisfies $0 < \alpha \leq 1$. (In Appendix A.1, the $\alpha \to 1$ limit is shown to be regular.) By solving $f(r_\pm) = 0$ w.r.t. μ and b^2, we obtain

$$\begin{aligned} \frac{h^5\mu}{L^3} &= -\frac{\beta_{7/2}}{\beta_1} + (\alpha^{3/2} + \alpha^{-3/2})h^2, \\ \frac{h^8 b^2}{L^6} &= \frac{12\beta_2}{\beta_1} - 12h^2, \end{aligned} \tag{3.10}$$

where $\beta_n \equiv \sum_{i=0}^{2n} \alpha^{i-n}$. With this parametrization, (2.15) and (3.9) become

$$\frac{2\pi - \delta_+}{2\pi - \delta_-} = \frac{\gamma_+ - 3\beta_1^2 h^2}{\gamma_- - 3\beta_1^2 h^2}\alpha^4 \tag{3.11}$$

and

$$\frac{\Phi^2}{(2\pi - \delta_+)(2\pi - \delta_-)} = \frac{16}{3}\frac{(\beta_2 - \beta_1 h^2)\beta_1^3}{(\gamma_+ - 3\beta_1^2 h^2)(\gamma_- - 3\beta_1^2 h^2)}, \tag{3.12}$$

respectively, where

$$\begin{aligned}\gamma_+ &= 3\alpha^3+6\alpha^2+9\alpha+12+15\alpha^{-1}+10\alpha^{-2}+5\alpha^{-3},\\ \gamma_- &= 5\alpha^3+10\alpha^2+15\alpha+12+9\alpha^{-1}+6\alpha^{-2}+3\alpha^{-3}.\end{aligned} \tag{3.13}$$

By using $\Delta\phi > 0$, $0 < \alpha \leq 1$ and $b^2 \geq 0$, it is shown that

$$0 < \frac{2\pi-\delta_+}{2\pi-\delta_-} \leq 1, \tag{3.14}$$

where the equality holds for $\alpha = 1$. The inequality (3.14) combined with the formula (2.11) implies that the tension of the brane at $r = r_-$ must be smaller than the tension of the brane at $r = r_+$. If this condition is not satisfied then it is expected that the cosmological constant on the brane cannot be non-negative or/and the geometry of the extra dimensions becomes dynamical. The warped volume Σ of extra dimensions defined in (3.7) is given by

$$\frac{\Sigma}{\sqrt{(2\pi-\delta_+)(2\pi-\delta_-)}} = \frac{2\beta_1^2}{3\sqrt{(\gamma_+-3\beta_1^2h^2)(\gamma_--3\beta_1^2h^2)}}. \tag{3.15}$$

3.2 Brane gravity on the IR brane

We now would like to see that the relation of the form (3.5) indeed holds at low energy on each brane. Since the warp factor r is smaller on the brane at $r = r_-$, we may call this brane an IR brane and the other brane at $r = r_+$ a UV brane. In this and the next subsections, we shall show the relation (3.5) on the IR and UV branes, respectively, by using numerical plots and determine the proportionality coefficient. In subsection 3.4 we show the same results analytically by low energy expansion.

Let us first consider the case where the phase transition takes place on the brane at $r = r_-$. In this case the deficit angle δ_+ around the other brane at $r = r_+$ and the flux Φ must be fixed. It is convenient to define the following dimensionless quantities:

$$\begin{aligned}\eta_- &\equiv \frac{2\pi-\delta_-}{2\pi-\delta_+},\\ h_- &\equiv L^2H_- = h\alpha^{-1/2},\\ \Phi_+ &\equiv \frac{|\Phi|}{2\pi-\delta_+} = \left[\frac{\Phi^2}{(2\pi-\delta_+)(2\pi-\delta_-)}\right]^{1/2}\eta_-^{1/2}\\ g_- &\equiv \frac{8\pi G_{N-}}{3}M_6^4L^2(2\pi-\delta_+) = \frac{1}{3}\left[\frac{\Sigma}{\sqrt{(2\pi-\delta_+)(2\pi-\delta_-)}}\right]^{-1}\eta_-^{-1/2}\alpha,\end{aligned} \tag{3.16}$$

where G_{N-} is defined by (3.6). The formulas (3.11), (3.12) and (3.15) give expressions of η_-, Φ_+^2 and g_- in terms of h_-^2 and α. The inequality (3.14) is written as $\eta_- \geq 1$, where the equality holds for $\alpha = 1$.

The quantities α, h_-, η_-, Φ_+ and g_- have the following physical meaning. The first quantity $\alpha \equiv r_-/r_+$ $(0 < \alpha \leq 1)$ is the ratio of the warp factor so that a small (or large) value of α corresponds to a large (or small, respectively) hierarchy. The second quantity h_- is the Hubble expansion rate on the brane at $r = r_-$ in the unit of the bulk curvature scale L^{-1}. Note that in this subsection, the phase transition is supposed to take place on this brane. Since the deficit angle δ_+ around the other brane at $r = r_+$ is determined by the tension of the other brane at $r = r_+$ (see the formula (2.11)) and has nothing to do with the phase transition on the brane at $r = r_-$, $2\pi - \delta_+$ is constant during the phase transition. Hence, the change of tension of the brane at $r = r_-$ is proportional to the change of the third quantity η_-. The proportionality coefficient depends on $2\pi - \delta_+$ but it does not change by the phase transition. Similarly, specifying the magnetic flux up to its signature is equivalent to specifying the forth quantity Φ_+. As already stated, all relevant equations are invariant under reflection of the sign of the flux and, thus, we do not need to keep track of the sign. The normalization of Φ_+ has been determined so that it does not change during the phase transition and that it absorbs the factor $(2\pi - \delta_+)^{-2}$ in the eq. (3.12) divided by (3.11). The final quantity g_- is proportional to the quantity G_{N-} which is defined by (3.6) and which shall be identified as effective Newton's constant on the brane at $r = r_-$. The normalization of g_- has been determined so that g_- is dimensionless and that it eliminates explicit dependence on $(2\pi - \delta_+)$ from eq. (3.19) below.

When the Hubble expansion rate on the brane vanishes, i.e. when $h^2 = 0$, the equations are simplified. Hereafter, a superscript (0) shows that the corresponding quantity is evaluated at $h^2 = 0$. In particular, it is easy to see that

$$\begin{aligned} \frac{1}{5} &\leq \Phi_+^{(0)2} < \infty, \\ \frac{10}{3} &\geq g_-^{(0)} > 0, \end{aligned} \tag{3.17}$$

where each quantity (with $h = 0$) is a monotonic function of α and the left and right values are values at $\alpha = 1$ and $\alpha = 0$, respectively.

Our aim in this paper is to show that, when a phase transition takes place on a brane, the Hubble expansion rate changes according to the standard Friedmann equation with Newton's constant given by the formula (3.6). In the present context where the phase transition takes place on the brane at $r = r_-$, we would like to see



Figure 1: Each line represents the relation between h_-^2 (vertical axis) and η_- (horizontal axis) for a fixed value of Φ_+^2 (= 0.3, 0.4, 0.5, 0.6, 0.7 from left to right). Note that the value of η_- is restricted to the region $\eta_- \geq 1$ as shown in (3.14). The domain shown in this figure can be joined to the domain shown in Fig. 3 by identifying the vertical line $\eta_- = 1$ in this figure to the vertical line $\eta_+ = 1$ in Fig. 3.

a relation between h_-^2 and η_- with Φ_+^2 fixed. Actually, by using equations (3.11) and (3.12), it is fairly easy to plot the curve $(\eta_-,\, h_-^2)$ for various fixed values of Φ_+^2. See Fig. 1. It is seen that the slope of each curve in Fig. 1 is negative near the horizontal axis, i.e. for small h_-^2. In other words, at low energy the Hubble expansion rate h_- indeed decreases as the brane tension σ_- and the deficit angle δ_- decrease (see the formula (2.11)) by a phase transition, and Newton's constant on the brane is positive. The intersection of each curve with the horizontal axis defines the value $\eta_-^{(0)}(\Phi_+^2)$ of η_- corresponding to $h_-^2 = 0$. Thus,

$$\sigma_-^{(0)}(\Phi_+^2) \equiv \left[2\pi - (2\pi - \delta_+)\eta_-^{(0)}(\Phi_+^2)\right] \cdot M^4 \tag{3.18}$$

is the critical value of tension σ_- for which the effective cosmological constant on the brane vanishes. If $\sigma_- > \sigma_-^{(0)}(\Phi_+^2)$ (or $\sigma_- < \sigma_-^{(0)}(\Phi_+^2)$) then the effective cosmological constant on the brane is positive (or negative, respectively).

The positivity of low energy effective Newton's constant is highlighted in Fig. 2, where the vertical axis is again h_-^2 but the horizontal axis is now $\eta_-^{(0)}(\Phi_+^2) - \eta_-$. The positive slope of the curve near the origin clearly defines low energy effective

Newton's constant since it determines how much the Hubble expansion rate changes as the brane tension changes. The dot lines in Fig. 2 are straight lines passing through the origin with the slope $g_-^{(0)}(\Phi_+^2)$, where $g_-^{(0)}(\Phi_+^2)$ is the value of g_- with $h_-^2 = 0$ for a given value of Φ_+^2. It is easily seen that the solid lines and the dot lines in Fig. 2 come in contact with each other at the origin. This fact implies that

$$h_-^2 = g_-^{(0)} \cdot (\eta_-^{(0)} - \eta_-) + O((\eta_-^{(0)} - \eta_-)^2). \tag{3.19}$$

By definition of g_- and the formula (2.11), this is equivalent to

$$H_-^2 = \frac{8\pi G_{N-}}{3}(\sigma_- - \sigma_-^{(0)}) + O((\sigma_- - \sigma_-^{(0)})^2), \tag{3.20}$$

which confirms that G_{N-} defined in (3.6) indeed has the meaning of 4-dimensional Newton's constant on the brane at $r = r_-$. The straight dot lines in Fig. 2 are good approximation to the solid lines in the regime where h_-^2 is sufficiently smaller than 1. This suggests that the Friedmann equation should be recovered at low energy where the Hubble expansion rate squared H_-^2 on the brane is sufficiently smaller than the bulk cosmological constant Λ_6 $(= 10L^{-2})$. In subsection 3.4, we shall show the relation (3.20) analytically.

3.3 Brane gravity on the UV brane

Next let us consider the case where a phase transition takes place on the UV brane at $r = r_+$. In this subsection, we shall show the relation (3.5) on the UV brane by using numerical plots and determine the proportionality coefficient. In the next subsection 3.4 we show the same results analytically by low energy expansion.

In the present case where a phase transition takes place on the brane at $r = r_+$, the deficit angle δ_- around the brane at $r = r_-$ and the flux Φ must be fixed. Similarly to the previous case, it is convenient to define the following dimensionless quantities:

$$\begin{aligned}
\eta_+ &\equiv \frac{2\pi - \delta_+}{2\pi - \delta_-}, \\
h_+ &\equiv LH_- = h\alpha^{1/2}, \\
\Phi_- &\equiv \frac{|\Phi|}{2\pi - \delta_-} = \left[\frac{\Phi^2}{(2\pi - \delta_+)(2\pi - \delta_-)}\right]^{1/2} \eta_+^{1/2} \\
g_+ &\equiv \frac{8\pi G_{N+}}{3} M_6^4 L^2 (2\pi - \delta_-) = \frac{1}{3}\left[\frac{\Sigma}{\sqrt{(2\pi - \delta_+)(2\pi - \delta_-)}}\right]^{-1} \eta_+^{-1/2}\alpha^{-1},
\end{aligned} \tag{3.21}$$

where G_{N+} is defined by (3.6). The formulas (3.11), (3.12) and (3.15) give expressions of η_+, Φ_-^2 and g_+ in terms of h_+^2 and α. The inequality (3.14) is written as $0 < \eta_+ \leq 1$, where the equality holds for $\alpha = 1$.



Figure 2: Solid lines are curves shown in Fig. 1 in a different coordinate. The vertical axis is again h_-^2 but the horizontal axis is now $\eta_-^{(0)}(\Phi_+^2) - \eta_-$, where $\eta_-^{(0)}(\Phi_+^2)$ is the value of η_- corresponding to $h_-^2 = 0$ for a given value of Φ_+^2. The value of Φ_+^2 for each curve is 0.3, 0.4, 0.5, 0.6, 0.7 from left to right. Dot lines represent straight lines passing through the origin with the slope $g_-^{(0)}(\Phi_+^2)$ for each value of Φ_+^2. These graphs confirm that G_{N-} defined by (3.6) indeed has the meaning of 4-dimensional Newton's constant on the brane at $r = r_-$.

The quantities h_+, η_+, Φ_- and g_+ have the physical meaning similar to h_-, η_-, Φ_+ and g_-, whose physical meaning has already been explained in the third paragraph of the previous subsection.

When the Hubble expansion rate on the brane vanishes, i.e. when $h^2 = 0$, the equations are simplified. As before, a superscript (0) shows that the corresponding quantity is evaluated at $h^2 = 0$. In particular, it is easy to see that

$$\begin{aligned} 0 &< \Phi_-^{(0)2} \leq \frac{1}{5}, \\ \infty &> g_+^{(0)} \geq \frac{10}{3} \end{aligned} \tag{3.22}$$

where each quantities (with $h = 0$) is a monotonic function of α and the left and right values are values at $\alpha = 0$ and $\alpha = 1$, respectively.

As already stated many times, our aim in this paper is to show that, when a phase transition takes place on a brane, the Hubble expansion rate changes according to the standard Friedmann equation with Newton's constant given by the formula (3.6). In this subsection we are considering a situation where the phase transition takes place on the brane at $r = r_+$. Thus, we would like to see a relation between h_+^2 and η_+ with Φ_-^2 fixed. Actually, by using equations (3.11) and (3.12), it is fairly easy to plot the curve $(\eta_+,\ h_+^2)$ for various fixed values of Φ_-^2. See Fig. 3. It is seen that the slope of each curve in Fig. 3 is negative near the horizontal axis, i.e. for small h_+^2. In other words, at low energy the Hubble expansion rate h_+ indeed decreases as the brane tension σ_+ and the deficit angle δ_+ decrease (see the formula (2.11)) by a phase transition, and Newton's constant on the brane is positive.

At first sight, this appears to contradict with the negativity of the slope of the curves in Fig. 1 near $h_-^2 = 0$: Fig. 3 indicates that η_+ decreases when h^2 increases from zero but Fig. 1 indicates that $\eta_- = 1/\eta_+$ also decreases when h^2 increases from zero. Actually, there is no contradiction. Along each curve in Fig. 1, what is fixed is not Φ_-^2 but Φ_+^2. If h^2 is increased from zero with Φ_-^2 fixed then Fig. 3 says that $\Phi_+^2 = \Phi_-^2/\eta_+^2$ should increase. In Fig. 1 this means that we have to move from a curve with the initial Φ_+^2 to different curves with larger Φ_+^2. Hence, if the value of η_- were the same then this would inevitably increase the value of h^2 from zero. Actually, since η_- is the reciprocal of η_+, Fig. 3 says that η_- should also increase. Therefore, when h^2 is increased from zero, there are two competitive effects which Fig. 3 implies: (i) the increase of Φ_+^2; (ii) the increase of η_-. The effect (i) tends to increase h^2 from zero but the effect (ii) tends to decrease h^2. The result of the competition is that the effect (i) wins. Hence, the negativity of the slopes of the curves both in Fig. 1 and Fig. 3 near the horizontal axis is consistent with each other.



Figure 3: Each line represents the relation between h_+^2 (vertical axis) and η_+ (horizontal axis) for a fixed value of Φ_-^2 (= 0.02, 0.08, 0.14, 0.2 from left to right). Note that the value of η_+ is restricted to the region $0 < \eta_+ \leq 1$ as shown in (3.14). For a relation between curves in this figure and Fig. 1, see the third-to-the-last paragraph in subsection 3.3. The domain shown in this figure can be joined to the domain shown in Fig. 1 by identifying the vertical line $\eta_+ = 1$ in this figure to the vertical line $\eta_- = 1$ in Fig. 1.



Figure 4: Solid lines are curves shown in Fig. 3 in a different coordinate. The vertical axis is again h_+^2 but the horizontal axis is now $\eta_+^{(0)}(\Phi_-^2) - \eta_+$, where $\eta_+^{(0)}(\Phi_-^2)$ is the value of η_+ corresponding to $h_+^2 = 0$ for a given value of Φ_-^2. The value of Φ_-^2 for each curve is 0.02, 0.08, 0.14, 0.2 from left to right. Dot lines represent straight lines passing through the origin with the slope $g_+^{(0)}(\Phi_-^2)$ for each value of Φ_-^2. These graphs confirm that G_{N+} defined by (3.6) indeed has the meaning of 4-dimensional Newton's constant on the brane at $r = r_+$.

The intersection of each curve in Fig. 3 with the horizontal axis defines the value $\eta_+^{(0)}(\Phi_-^2)$ of η_+ corresponding to $h_+^2 = 0$. Thus,

$$\sigma_+^{(0)}(\Phi_-^2) \equiv \left[2\pi - (2\pi - \delta_-)\eta_+^{(0)}(\Phi_-^2)\right] \cdot M^4 \tag{3.23}$$

is the critical value of tension σ_+ for which the effective cosmological constant on the brane vanishes. If $\sigma_+ > \sigma_+^{(0)}(\Phi_-^2)$ (or $\sigma_+ < \sigma_+^{(0)}(\Phi_-^2)$) then the effective cosmological constant on the brane is positive (or negative, respectively).

The positivity of low energy effective Newton's constant is highlighted in Fig. 4, where the vertical axis is again h_+^2 but the horizontal axis is now $\eta_+^{(0)}(\Phi_-^2) - \eta_+$. The positive slope of the curve near the origin clearly defines low energy effective Newton's constant since it determines how much the Hubble expansion rate changes as the brane tension changes. The dot lines in Fig. 4 are straight lines passing through the origin with the slope $g_+^{(0)}(\Phi_-^2)$, where $g_+^{(0)}(\Phi_-^2)$ is the value of g_+ with $h_+^2 = 0$ for a given value of Φ_-^2. It is easily seen that the solid lines and the dot lines in Fig. 4

come in contact with each other at the origin. This fact implies that

$$h_+^2 = g_+^{(0)} \cdot (\eta_+^{(0)} - \eta_+) + O((\eta_+^{(0)} - \eta_+)^2). \tag{3.24}$$

By definition of g_+ and the formula (2.11), this is equivalent to

$$H_+^2 = \frac{8\pi G_{N+}}{3}(\sigma_+ - \sigma_+^{(0)}) + O((\sigma_+ - \sigma_+^{(0)})^2), \tag{3.25}$$

which confirms that G_{N+} defined in (3.6) indeed has the meaning of 4-dimensional Newton's constant on the brane at $r = r_+$. The straight dot lines in Fig. 4 are good approximation to the solid lines in the regime where h_+^2 is sufficiently smaller than 1. This suggests that the Friedmann equation should be recovered at low energy where the Hubble expansion rate squared H_+^2 on the brane is sufficiently smaller than the bulk cosmological constant Λ_6 $(= 10L^{-2})$. In the next subsection 3.4, we shall show the relation (3.25) analytically.

3.4 Low energy expansion

In this subsection we show the low-energy relations (3.20) and (3.25) analytically, where Newton's constant $G_{N\pm}$ is given by (3.6). For this purpose we shall perform perturbative expansion of relevant quantities and equations w.r.t. h^2 $(= h_+h_-)$ and show that (3.19) and (3.24) indeed hold in the lowest order in the perturbative expansion.

Let us first consider the case considered in subsection 3.2, where the phase transition takes place on the brane at $r = r_-$. In this case we expand α and η_- as

$$\alpha = \sum_{n=0}^{\infty} \alpha_-^{(n)}(\Phi_+^2)h^{2n}, \quad \eta_- = \sum_{n=0}^{\infty} \eta_-^{(n)}(\Phi_+^2)h^{2n}. \tag{3.26}$$

Since Φ_+ is fixed during the phase transition, we formally consider Φ_+ as a quantity of order $O(h^0)$. It is straightforward to expand (3.12) and (3.11) w.r.t. h^2. In the order $O(h^0)$, we obtain

$$\begin{aligned} \Phi_+^2 &= \frac{16\beta_2^{(0)}\beta_1^{(0)3}}{3\gamma_+^{(0)2}\alpha_-^{(0)4}}, \\ \eta_-^{(0)} &= \frac{\gamma_-^{(0)}}{\gamma_+^{(0)}\alpha_-^{(0)4}}, \end{aligned} \tag{3.27}$$

where $\beta_n^{(0)}$ and $\gamma_\pm^{(0)}$ are β_n and $\gamma_\pm$, respectively, with α replaced by $\alpha_-^{(0)}$. On the other hand, the zeroth order part of eq. (3.15) gives the value $g_-^{(0)}(\Phi_+^2)$ of g_- with $h_-^2 = 0$

for a given value of Φ_+^2 as

$$g_-^{(0)} = \frac{\gamma_+^{(0)}\alpha_-^{(0)3}}{2\beta_1^{(0)2}}. \tag{3.28}$$

In the order $O(h^2)$, eq. (3.12) divided by (3.11) gives an expression of $\alpha_-^{(1)}$ in terms of $\alpha_-^{(0)}$ as

$$\frac{\alpha_-^{(1)}}{\alpha_-^{(0)}} = \frac{\beta_1^{(0)2}A_-^{(0)}}{B_-^{(0)}}, \tag{3.29}$$

where

$$\begin{aligned} A_-^{(0)} &= 3\alpha_-^{(0)3} + 6\alpha_-^{(0)2} + 9\alpha_-^{(0)} + 6 + 3\alpha_-^{(0)-1} + 2\alpha_-^{(0)-2} + \alpha_-^{(0)-3}, \\ B_-^{(0)} &= 15\alpha_-^{(0)6} + 60\alpha_-^{(0)5} + 150\alpha_-^{(0)4} + 258\alpha_-^{(0)3} + 357\alpha_-^{(0)2} + 430\alpha_-^{(0)} + 460 \\ &\quad +430\alpha_-^{(0)-1} + 357\alpha_-^{(0)-2} + 258\alpha_-^{(0)-3} + 150\alpha_-^{(0)-4} + 60\alpha_-^{(0)-5} + 15\alpha_-^{(0)-6}. \end{aligned} \tag{3.30}$$

Here, as already explained, Φ_+ was considered as a quantity of order $O(h^0)$ in the derivation of the above expression of $\alpha_-^{(0)}$. On the other hand, the reciprocal of eq. (3.11) in the order $O(h^2)$ gives

$$\eta_-^{(1)} = -\frac{2\beta_1^{(0)2}}{\gamma_+^{(0)}\alpha_-^{(0)4}} = -\frac{1}{g_-^{(0)}\alpha_-^{(0)}}. \tag{3.31}$$

Now, the definition of $\eta_-^{(0,1)}$

$$\eta_- = \eta_-^{(0)} + \eta_-^{(1)}h^2 + O(h^4) \tag{3.32}$$

is solved w.r.t. h^2 as

$$h^2 = \frac{\eta_- - \eta_-^{(0)}}{\eta_-^{(1)}} + O(h^4), \tag{3.33}$$

or equivalently

$$h_-^2 = \frac{h^2}{\alpha} = g_-^{(0)} \cdot \left(\eta_-^{(0)} - \eta_-\right) + O(h_-^4). \tag{3.34}$$

This proves (3.19). By the definition of g_- and the formula (2.11), this is equivalent to (3.20).

It is easy to perform a similar analysis for the other case considered in subsection 3.3, where the phase transition takes place on the brane at $r = r_+$, and to derive (3.19) and (3.25) analytically. In this case we expand α and η_+ as

$$\alpha = \sum_{n=0}^{\infty} \alpha_+^{(n)}(\Phi_-^2)h^{2n}, \quad \eta_+ = \sum_{n=0}^{\infty} \eta_+^{(n)}(\Phi_-^2)h^{2n}. \tag{3.35}$$

Since Φ_- is fixed during the phase transition, we formally consider Φ_- as a quantity of order $O(h^0)$. It is straightforward to expand (3.12) and (3.11) w.r.t. h^2. In the order $O(h^0)$, we obtain

$$\begin{aligned} \Phi_-^2 &= \frac{16\beta_2^{(0)}\beta_1^{(0)3}\alpha_+^{(0)4}}{3\gamma_-^{(0)2}}, \\ \eta_+^{(0)} &= \frac{\gamma_+^{(0)}\alpha_+^{(0)4}}{\gamma_-^{(0)}}, \end{aligned} \tag{3.36}$$

where $\beta_n^{(0)}$ and $\gamma_\pm^{(0)}$ are β_n and $\gamma_\pm$, respectively, with α replaced by $\alpha_+^{(0)}$. On the other hand, the zeroth order part of eq. (3.15) gives the value $g_+^{(0)}(\Phi_+^2)$ of g_+ with $h_+^2 = 0$ for a given value of Φ_-^2 as

$$g_+^{(0)} = \frac{\gamma_-^{(0)}}{2\beta_1^{(0)2}\alpha_+^{(0)3}}. \tag{3.37}$$

In the order $O(h^2)$, eq. (3.12) multiplied by (3.11) gives an expression of $\alpha_+^{(1)}$ in terms of $\alpha_+^{(0)}$ as

$$\frac{\alpha_+^{(1)}}{\alpha_+^{(0)}} = \frac{\beta_1^{(0)2}A_+^{(0)}}{B_+^{(0)}}, \tag{3.38}$$

where

$$\begin{aligned} A_+^{(0)} &= 3\alpha_+^{(0)-3} + 6\alpha_+^{(0)-2} + 9\alpha_+^{(0)-1} + 6 + 3\alpha_+^{(0)} + 2\alpha_+^{(0)2} + \alpha_+^{(0)3}, \\ B_+^{(0)} &= 15\alpha_+^{(0)-6} + 60\alpha_+^{(0)-5} + 150\alpha_+^{(0)-4} + 258\alpha_+^{(0)-3} + 357\alpha_+^{(0)-2} + 430\alpha_+^{(0)-1} + 460 \\ &\quad +430\alpha_+^{(0)} + 357\alpha_+^{(0)2} + 258\alpha_+^{(0)3} + 150\alpha_+^{(0)4} + 60\alpha_+^{(0)5} + 15\alpha_+^{(0)6}. \end{aligned} \tag{3.39}$$

On the other hand, eq. (3.11) in the order $O(h^2)$ gives

$$\eta_+^{(1)} = -\frac{2\beta_1^{(0)2}\alpha_+^{(0)4}}{\gamma_-^{(0)}} = -\frac{\alpha_+^{(0)}}{g_+^{(0)}}. \tag{3.40}$$

Now, the definition of $\eta_+^{(0,1)}$

$$\eta_+ = \eta_+^{(0)} + \eta_+^{(1)}h^2 + O(h^4) \tag{3.41}$$

is solved w.r.t. h^2 as

$$h^2 = \frac{\eta_+ - \eta_+^{(0)}}{\eta_+^{(1)}} + O(h^4), \tag{3.42}$$

or equivalently

$$h_+^2 = h^2\alpha = g_+^{(0)} \cdot \left(\eta_+^{(0)} - \eta_+\right) + O(h_+^4). \tag{3.43}$$

This proves (3.24). By the definition of g_+ and the formula (2.11), this is equivalent to (3.25).

It is straightforward to extend the analysis in this subsection to to higher oder in the expansion w.r.t. h^2 and to interpret the result as higher-order corrections to the effective Friedmann equation. In Appendix A.3 the results in the order $O(h^4)$ is summarized and it is shown that higher order corrections can be ignored when the Hubble expansion rate $H_\pm$ on a brane is sufficiently lower than the bulk curvature scale L^{-1}.

4 Summary and discussion

We have considered a 6-dimensional model of warped flux compactification and analyzed gravity sourced by a 3-brane in a simplified setup. This setup includes a warped geometry, compactification, a flux, and one or two 3-brane(s). We have considered a situation where the tension of a 3-brane changes by a phase transition on the brane. Assuming that the tension is almost constant in deep inside the old and new phases, we have investigated the relation between the change of tension and the change of the Hubble expansion rate on the brane. The relation is the same as that inferred by the 4-dimensional Einstein equation, provided that Newton's constant is given by the formula (3.6).

We have explicitly seen how the induced geometry changes according to a change of brane source. With the static ansatz for the bulk, the bulk geometry is uniquely determined by the brane tensions and the value of the conserved flux. As a consequence, the Hubble expansion rate induced on the brane is determined. Provided that the solutions considered in this paper are dynamically stable [54], this result implies that, when a brane tension changes, the bulk geometry should evolve toward the corresponding unique configuration. This process determines the evolution of the bulk geometry and, thus, the induced geometry on the brane. In this way, the induced geometry responses to the brane source.

Although the situation we have considered is rather restrictive, the physical picture we have obtained seems more general. Provided that all moduli are stabilized, when a brane source changes slowly compared to the time scale of the moduli dynamics, the bulk geometry should quickly settle to a configuration which is determined by the boundary condition, i.e. the brane source(s), values of conserved quantities and the regularity of the other region of the extra dimensions. As a consequence of the change of the bulk geometry, the induced geometry on the brane responses to the evolution of the matter source on the brane.

Suppose that all moduli are stabilized in a $(4+n)$-dimensional warped compactification. From the viewpoint of the effective field theory with massive moduli integrated out, it is in general expected that the Einstein gravity should be recovered at low energy [5] and that 4-dimensional Newton's constant should be given by the formula (3.3):

$$\frac{1}{8\pi G_N} = (M_{4+n})^{2+n} \int d^n y \sqrt{\gamma} \left[\frac{r(y)}{r(y_0)} \right]^2 . \tag{4.1}$$

(See the first paragraph of Sec. 3 for the reasoning leading to this expectation.) Here, M_{4+n} is the $(4+n)$-dimensional Planck mass, $r^2(y)$ is the warp factor of mass dimension -2 depending on the coordinates of compact extra dimensions y^i, $\gamma_{ij}dy^i dy^j$ is the metric of the extra dimensions (see eq. (3.1)) and the brane source is supposed to be localized at $y^i = y_0^i$. Although the effective field theory approach is usually a powerful and useful tool to obtain a correct answer quickly, the result must be checked by independent methods. We expect that this formula should hold at low energy in a wide class of warped compactification if all moduli are stabilized and made massive.

The formula (3.3) was already proved to be correct in a Randall-Sundrum type, codimension 1 braneworlds with radion stabilization [37, 38, 39, 40, 41, 42, 43] [6]. The result of the present paper confirms the same formula for codimension 2 braneworlds in a simplified situation.

Future subjects include the analysis of dynamical stability [54], the recovery of the 4-dimensional linearized Einstein equation for inhomogeneous perturbations, evolution of FRW universe on the brane, extensions to systems with a dilaton, and so on.

The simple 6-dimensional setup in this paper at the very least provides non-trivial evidence for validity of the expected formula (3.3) for effective Newton's constant in warped flux compactification. Thus, let us now briefly discuss application of the formula (3.3) to cosmology with type IIB warped flux compactification considered in refs. [1, 14, 15, 16, 17, 18]. The 10-dimensional geometry is of the form

$$ds_{10}^2 = h^{-1/2}(\tau) g^{(4)}_{\mu\nu} dx^\mu dx^\nu + h^{1/2}(\tau) \gamma_{mn} d\psi^m d\psi^n, \tag{4.2}$$

---

[5] The stress-energy tensor in the 4-dimensional effective Einstein equation must be calculated from a 4-dimensional effective action after integrating out all massive moduli. The effective stress-energy tensor defined in this way is not necessarily the same as the surface stress-energy tensor (the so called Lanczos tensor [55] in Israel's junction condition [50] for codimension 1 cases). If moduli are not stabilized, it is a non-trivial issue whether a 4-dimensional theory is recovered. See eg. [56, 57] for this issue in the context of Randall-Sundrum type braneworlds.

[6] See the footnote after (3.3).

where γ_{mn} is the metric of a 6-dimensional compact geometry (Calabi-Yau space) depending on the coordinates ψ^m ($m = 5, \cdots, 10$) of extra dimensions, $h(\tau)$ is a function of the radial coordinate τ ($\in \{\psi^m\}$) of the compact geometry, and $g^{(4)}_{\mu\nu}$ is a 4-dimensional metric. It is supposed that the function $h(\tau)$ is of order 1 in the bulk of the compact geometry but that there is a region called a warped throat where $h(\tau)$ becomes exponentially large. In the warped throat region the warp factor $h^{-1/2}(\tau)$ becomes exponentially small.

In most of refs. [14, 15, 16, 17, 18] it was (implicitly) assumed that what drives (or at least affects) the 4-dimensional cosmology is a brane in the throat region but that our 4-dimensional world is somewhere in the Calabi-Yau space where the warp factor $h^{-1/2}(\tau)$ is of order unity. In other words, the physical metric of our 4-dimensional world is $g^{(4)}_{\mu\nu}$ up to a normalization constant of order unity, while the induced metric on the brane is not $g^{(4)}_{\mu\nu}$ but

$$g^{(b)}_{\mu\nu} = h^{-1/2}(\tau_0) \cdot g^{(4)}_{\mu\nu}, \tag{4.3}$$

where τ_0 is the value of the radial coordinate τ at the position of the brane. At first sight, it might seem non-trivial how the brane at $\tau = \tau_0$ drives the cosmology of our 4-dimensional world $g^{(4)}_{\mu\nu}$ since the warp factor $h^{-1/2}(\tau_0)$ is exponentially small. The approach adopted in the literature is to write down the brane action in terms of $g^{(4)}_{\mu\nu}$ and couple it to the Einstein gravity in the frame $g^{(4)}_{\mu\nu}$, assuming that Newton's constant $G^{(4)}_N$ is given by the Kaluza-Klein like formula

$$\frac{1}{8\pi G^{(4)}_N} = M^8_{10} V_6, \tag{4.4}$$

where V_6 is the volume of the Calabi-Yau space.

We can justify this approach by using the formula (3.3). First, the induced geometry on the brane is, as already stated, the conformally transformed metric $g^{(b)}_{\mu\nu}$ given in (4.3). Thus, the stress energy tensor $T^{(b)}_{\mu\nu}$ in the brane frame $g^{(b)}_{\mu\nu}$ and the stress energy tensor $T^{(4)}_{\mu\nu}$ in the frame $g^{(4)}_{\mu\nu}$ are related to each other as

$$T^{(b)\nu}_{\mu} = h(\tau_0) \cdot T^{(4)\nu}_{\mu}, \tag{4.5}$$

where τ_0 is again the value of the radial coordinate τ at the position of the brane and it has been assumed that the value of $h(\tau)$ at the position of our 4-dimensional world is normalized to 1. Now, the formula (3.3) applied to this setup says that we can use the effective Einstein equation of the form

$$G^{(b)\nu}_{\mu} = 8\pi G^{(b)}_N T^{(b)\nu}_{\mu}, \tag{4.6}$$

where $G_{\mu}^{(b)\nu}$ is the Einstein tensor of the brane metric $g_{\mu\nu}^{(b)}$ and Newton's constant $G_N^{(b)}$ on the brane is given by

$$\frac{1}{8\pi G_N^{(b)}} = M_{10}^8 \int d^6\psi h^{3/2}(\tau)\sqrt{\gamma} \cdot \left(\frac{h(\tau)}{h(\tau_0)}\right)^{-1/2}. \tag{4.7}$$

By the assumption that the function $h(\tau)$ is of order unity in the bulk of extra dimensions, this is reduced to

$$\frac{1}{8\pi G_N^{(b)}} \simeq h^{1/2}(\tau_0) M_{10}^8 V_6 \simeq \frac{h^{1/2}(\tau_0)}{8\pi G_N^{(4)}}, \tag{4.8}$$

where V_6 is again the volume of the Calabi-Yau space. This result implies that the effective Einstein equation on the brane (4.6) is equivalent to

$$G_{\mu}^{(4)\nu} = 8\pi G_N^{(4)} T_{\mu}^{(4)\nu}, \tag{4.9}$$

where we have used the conformal transformation

$$G_{\mu}^{(b)\nu} = h^{1/2}(\tau_0) G_{\mu}^{(4)\nu} \tag{4.10}$$

between Einstein tensors $G_{\mu}^{(b)\nu}$ and $G_{\mu}^{(4)\nu}$ of $g_{\mu\nu}^{(b)}$ and $g_{\mu\nu}^{(4)}$, respectively. Therefore, we have justified the approach adopted in the literature by using the formula (3.3). The 4-dimensional fields resulting from the brane action indeed couple to the 4-dimensional metric $g_{\mu\nu}^{(4)}$ with Newton's constant $G_N^{(4)}$ given by the Kaluza-Klein like formula (4.4).

Acknowledgements

We would like to thank Andrei Frolov, Yoshiaki Himemoto, Lev Kofman, Kazuya Koyama, Hideaki Kudoh, Tetsuya Shiromizu, Jiro Soda, Keitaro Takahashi and Takahiro Tanaka for useful discussions and/or comments. We are grateful to Katsuhiko Sato for continuing encouragement. Y. S. and H. Y. are supported by JSPS.

Appendices

A.1 The $\alpha \to 1$ limit

In the limit $\alpha \equiv r_-/r_+ \to 1$, the coordinate distance between two branes vanishes and, thus, the bulk geometry appears to collapse. However, as we shall see below, the proper distance between $r = r_-$ and r_+ does not vanish and the geometry of extra dimensions remains regular.

When α is sufficiently close to 1, the function $f(r)$ between $r_\pm$ is approximated by

$$f(r) \simeq a^2 - b^2(r - r_0)^2, \tag{A.1}$$

where r_0, a and b are positive constants and the $\alpha \to 1$ limit corresponds to sending $a \to +0$ with b finite. The bulk coordinate r is restricted to the interval $-a/b \leq r - r_0 \leq a/b$. With the new coordinate system $(\theta,\, \varphi)$ defined by

$$\theta = \cos^{-1}\left[\frac{b}{a}(r - r_0)\right], \quad \varphi = ab\phi, \tag{A.2}$$

the metric of the extra dimension becomes that of a round sphere with radius $1/b$:

$$\frac{dr^2}{f(r)} + f(r)d\phi^2 \simeq \frac{1}{b^2}(d\theta^2 + \sin^2\theta d\varphi^2). \tag{A.3}$$

Evidently, the coordinate θ runs over the full interval $[0, \pi]$. On the other hand, the period of the coordinate φ appears to collapse since the coefficient ab in the definition (A.2) of φ vanishes in the $\alpha \to 1$ limit. Actually, this is not the case. The "surface gravity" $\kappa_\pm$ defined in (2.13) is $\kappa_+ \simeq \kappa_- \simeq ab$ and the period of the old coordinate is, thus, $\Delta\phi \simeq (2\pi - \delta_\pm)/(ab)$. Therefore, the new coordinate φ has the period

$$\Delta\varphi = ab\Delta\phi \simeq 2\pi - \delta_+ \simeq 2\pi - \delta_-, \tag{A.4}$$

which is indeed finite. Thus, the geometry of extra dimensions is nothing but a round sphere with a deficit angle $\delta_+ \simeq \delta_-$, i.e. a football-shaped extra-dimensions considered in [45, 46, 47]. In those papers a Z_2 symmetry is assumed so that tensions of two 3-branes are identical. With the Z_2 symmetry, $\eta_\pm$ is always 1 but $\Phi_\pm = \Phi/(2\pi - \delta_\pm)$ changes when the brane tension changes. Accordingly, as can be seen from Figs. 1 and 3, the Hubble expansion rate on the brane changes when the brane tension changes.

The warp factor r becomes constant in the $\alpha \to 1$ limit and, thus, the 6-dimensional solution in the $\alpha \to 1$ limit corresponds to an unwarped flux compactification of the type considered by Arkani-Hamed, Dimopoulos and Dvali [44].

A.2 The $h \to 0$ limit

The limit $h \to 0$ appears to be singular, but it is not. Indeed, by the coordinate change

$$r \to \frac{\bar{r}}{h}, \quad \phi \to h\bar{\phi}, \quad ds_4^2 \to h^2 d\bar{s}_4^2, \quad \mu \to \frac{\bar{\mu}}{h^5}, \quad b \to \frac{\bar{b}}{h^4}, \tag{A.5}$$

the $h \to 0$ limit of the bulk solution becomes a locally regular form:

$$\begin{aligned} ds_6^2 &= \bar{r}^2 d\bar{s}_4^2 + \frac{d\bar{r}^2}{\bar{f}(\bar{r})} + \bar{f}(\bar{r}) d\bar{\phi}^2, \\ A_M dx^M &= \frac{\bar{b}}{3\bar{r}^3} d\bar{\phi}, \end{aligned} \tag{A.6}$$

where

$$\bar{f}(\bar{r}) = -\frac{\Lambda_6}{10}\bar{r}^2 - \frac{\bar{\mu}}{\bar{r}^3} - \frac{\bar{b}^2}{12\bar{r}^6}, \tag{A.7}$$

and $d\bar{s}_4^2$ is the metric of the 4-dimensional Minkowski spacetime. This configuration of course satisfies the 6-dimensional Einstein equation and is related by a double Wick rotation to a topological black hole with the horizon topology $\mathbf{R}^4$.

The new coordinate $\bar{\phi}$ is identified as

$$\bar{\kappa}_\pm \bar{\phi} \sim \bar{\kappa}_\pm \bar{\phi} + (2\pi - \delta_\pm), \tag{A.8}$$

where

$$\bar{\kappa}_\pm \equiv \mp \frac{1}{2} \bar{f}'(\bar{r}_\pm). \tag{A.9}$$

This means that the period of $\bar{\phi}$ is finite if the brane tensions are finite. Therefore, the geometry remains regular in the $h \to 0$ limit.

A.3 Higher-order correction

It is straightforward to extend the analysis in subsection 3.4 to higher order in the expansion w.r.t. h^2 and to interpret the result as higher-order corrections to the effective Friedmann equation. In this appendix we just summarize the result up to the order $O(h^4)$.

The result up to the order $O(h^4)$ is summarized as

$$\left[1 - \frac{h_\pm^2}{h_{*\pm}^2} + O(h_\pm^4)\right] h_\pm^2 = g_\pm^{(0)} \cdot \left(\eta_\pm^{(0)} - \eta_\pm\right), \tag{A.10}$$

where

$$h_{*\pm}^2 = \frac{B_\pm^{(0)}}{\beta_1^2 A_\pm^{(0)}} \alpha_\pm^{(0)\pm 1}. \tag{A.11}$$

This is equivalent to

$$\left[1 - \frac{H_\pm^2}{H_{*\pm}^2} + O\left((H_\pm^2/\Lambda_6)^2\right)\right] H_\pm^2 = \frac{8\pi G_{N\pm}}{3}(\sigma_\pm - \sigma_\pm^{(0)}), \tag{A.12}$$

where $H_{*\pm}^2 = h_{*\pm}^2/L^2$, and includes a higher order correction to the effective Friedmann equation suppressed by $H_{*\pm}$. It is easy to show that

$$5 < L^2 H_{*+}^2 \le \left(\frac{10}{3}\right)^2, \quad \left(\frac{10}{3}\right)^2 \le L^2 H_{*-}^2 < \infty. \tag{A.13}$$

This means that higher order corrections can be ignored when the Hubble expansion rate $H_\pm$ on a brane is sufficiently lower than the bulk curvature scale L^{-1}.

References

[1] S. Kachru, R. Kallosh, A. Linde and S. P. Trivedi, Phys. Rev. **D68**, 046005 (2003) [hep-th/0301240].

[2] C. Escoda, M. Gomez-Reino and F. Quevedo, JHEP **0311**, 065 (2003) [hep-th/0307160].

[3] C. P. Burgess, R. Kallosh and F. Quevedo JHEP **0310**, 056 (2003) [hep-th/0309187].

[4] E. Silverstein, "(A)dS Backgrounds from Asymmetric Orientifolds", hep-th/0106209; A. Maloney, E. Silverstein and A. Strominger, "de Sitter Space in Noncritical String Theory", hep-th/0205316.

[5] M. Becker, G. Curio and A. Krause, Nucl. Phys. **B693**, 223 (2004) [hep-th/0403027].

[6] P. K. Townsend and M. N. R. Wohlfarth, Phys. Rev. Lett. **91**, 061302 (2003) [hep-th/0303097].

[7] N. Ohta, Phys. Rev. Lett. **91**, 061303 (2003) [hep-th/0303238]; Prog. Theor. Phys. **110**, 269 (2003) [hep-th/0304172]; Int. J. Mod. Phys. **A20**, 1 (2005) [hep-th/0411230].

[8] S. Roy, "Accelerating cosmologies from M/String theory compactifications", Phys. Lett. **B567**, 322 (2003) [hep-th/0304084].

[9] I. P. Neupane and D. L. Wiltshire, "Accelerating cosmologies from compactification with a twist", hep-th/0502003; "Cosmic acceleration from M-theory on twisted spaces", hep-th/0504135.

[10] K. Becker, M. Becker and A. Krause, Nucl. Phys. **B715**, 349 (2005) [hep-th/0501130].

[11] I. R. Klebanov and M. J. Strassler, JHEP **0008**, 052 (2000) [hep-th/0007191].

[12] S. B. Giddings, S. Kachru, J. Polchinski, Phys. Rev. **D66**, 106006 (2002) [hep-th/0105097].

[13] E. Witten, Nucl. Phys. **B474**, 343 (1996) [hep-th/9604030].

[14] S. Kachru, R. Kallosh, A. Linde, J. Maldacena, L. McAllister, S. P. Trivedi, JCAP **0310**, 013 (2003) [hep-th/0308055].

[15] J. J. Blanco-Pillado, C. P. Burgess, J. M. Cline, C. Escoda, M. Gomez-Reino, R. Kallosh, A. Linde and F. Quevedo, JHEP **0411**, 063 (2004) [hep-th/0406230].

[16] A. Buchel and Radu Roiban, Phys. Lett. **B590**, 284 (2004) [hep-th/0311154]; M. Berg and M. Haack, Phys. Rev. **D71**, 026005 (2005) [hep-th/0404087]; C. P. Burgess, J. M. Cline, H. Stoica and F. Quevedo, JHEP **0409**, 033 (2004) [hep-th/0403119]; O. DeWolfe, S. Kachru and H. Verlinde, JHEP **0405**, 017 (2004) [hep-th/0403123]; N. Iizuka and S. P. Trivedi, Phys. Rev. **D70**, 043519 (2004) [hep-th/0403203]; A. Buchel and A. Ghodsi, Phys. Rev. **D70**, 126008 (2004) [hep-th/0404151]; H. Firouzjahi and S. H. Tye, JCAP **0503**, 009 (2005) [hep-th/0501099]; J. P. Hsu, R. Kallosh and S. Prokushkin, JCAP **0312**, 009 (2003) [hep-th/0311077]; H. Firouzjahi and S. H. Tye, Phys. Lett. **B584**, 147 (2004) [hep-th/0312020]; J. P. Hsu and R. Kallosh, JHEP **0404**, 042 (2004) [hep-th/0402047]; F. Koyama, Y. Tachikawa and T. Watari, Phys. Rev. **D69**, 106001 (2004), Erratum-ibid. **D70**, 129907 (2004) [hep-th/0311191]; L. McAllister, "An Inflaton Mass Problem in String Inflation from Threshold Corrections to Volume Stabilization", hep-th/0502001; K. Dasgupta, J. P. Hsu, R. Kallosh, A. Linde and M. Zagermann, JHEP **0408**, 030 (2004) [hep-th/0405247]; P. Chen, K. Dasgupta, K. Narayan, M. Shmakova and M. Zagermann, "Brane inflation, solitons and cosmological solutions: I", hep-th/0501185; Yun-Song Piao, "D/antiD Dark Energy in String Warped Compactification", gr-qc/0502006; D. Cremades, F. Quevedo and A. Sinha, "Warped Tachyonic Inflation in Type IIB Flux Compactifications and the Open-string Completeness Conjecture", hep-th/0505252.

[17] J. Polchinski, "Introduction to Cosmic F- and D-strings", hep-th/0412244 and references therein.

[18] S. Mukohyama, "$\bar{D}$-brane as Dark Matter in Warped String Compactification", hep-th/0505042.

[19] G. R. Dvali and S. H. Tye, Phys. Lett. **B450**, 72 (1999) [hep-ph/9812483].

[20] F. Quevedo, Class. Quant. Grav. **19**, 5721 (2002) [hep-th/0210292] and references therein.

[21] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999) [hep-th/9906064].

[22] J. Garriga and T. Tanaka, Phys. Rev. Lett. **84**, 2778 (2000) [hep-th/9911055].

[23] M. Sasaki, T. Shiromizu and K. Maeda, Phys. Rev. **D62**, 024008 (2000) [hep-th/9912233].

[24] S. B. Giddings, E. Katz, and L. Randall, J. High Energy Phys. **03**, 023 (2000) [hep-th/0002091].

[25] A. Karch and L. Randall, JHEP **0105**, 008 (2001) [hep-th/0011156].

[26] T. Shiromizu, K. Maeda and M. Sasaki, Phys. Rev. **D62**, 024012 (2000) [gr-qc/9910076].

[27] C. Csaki, M. Graesser, C. F. Kolda and J. Terning, Phys. Lett. **B462**, 34 (1999) [hep-ph/9906513].

[28] J. M. Cline, C. Grojean and G. Servant, Phys. Rev. Lett. **83** 4245 (1999) [hep-ph/9906523].

[29] E. E. Flanagan, S. H. H. Tye, I. Wasserman, Phys. Rev. **D62**, 044039 (2000) [hep-ph/9910498].

[30] P. Binétruy, C. Deffayet, U. Ellwanger and D. Langlois, Phys. Lett. **B477**, 285 (2000) [hep-th/9910219].

[31] S. Mukohyama, Phys. Lett. **B473**, 241 (2000) [hep-th/9911165].

[32] P. Kraus, JHEP **9912**, 011 (1999) [hep-th/9910149].

[33] D. Ida, JHEP **0009**, 014 (2000) [gr-qc/9912002].

[34] S. Mukohyama, T. Shiromizu and K. Maeda, Phys. Rev. **D62**, 024028 (2000), Erratum-ibid. **D63**, 029901 (2001) [hep-th/9912287].

[35] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [hep-ph/9905221].

[36] W. D. Goldberger and M. B. Wise, Phys. Rev. Lett. **83**, 4922 (1999) [hep-th/0007065].

[37] T. Tanaka and X. Montes, Nucl. Phys. **B582**, 259 (2000) [hep-th/0001092].

[38] C. Csaki, M. L. Graesser and G. D. Kribs, Phys. Rev. **D63**, 065002 (2001) [hep-th/0008151].

[39] S. Mukohyama and L. Kofman, Phys. Rev. **D65**, 124025 (2002) [hep-th/0112115].

[40] S. Mukohyama, Phys. Rev. **D65**, 084036 (2002) [hep-th/0112205].

[41] H. Kudoh and T. Tanaka, Phys. Rev. **D65**, 104034 (2002) [hep-th/0112013]; Phys. Rev. **D67**, 044011 (2003) [hep-th/0205041].

[42] C. Csaki, M. Graesser and L. Randall, Phys. Rev. **D62**, 045015 (2000) [hep-ph/9911406].

[43] P. Kanti, I. I. Kogan, K. A. Olive and M. Pospelov, Phys. Lett. **B468**, 31 (1999) [hep-ph/9909481]; P. Kanti, K. A. Olive and M. Pospelov, Phys. Rev. **D62**, 126004 (2000) [hep-ph/0005146].

[44] N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. **B429**, 263 (1998) [hep-ph/9803315]; I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. **B436**, 257 (1998) [hep-ph/9804398]; N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Rev. **D59**, 086004 (1999) [hep-ph/9807344].

[45] S. M. Carroll and M. M. Guica, "Sidestepping the Cosmological Constant with Football Shaped Extra Dimensions", hep-th/0302067.

[46] J. Garriga and M. Porrati, JHEP **0408**, 028 (2004) [hep-th/0406158].

[47] I. Navarro, Class. Quant. Grav. **20**, 3603 (2003) [hep-th/0305014]; JCAP **0309**, 004 (2003) [hep-th/0302129].

[48] J. Vinet and J. M. Cline, Phys. Rev.**D71**, 064011 (2005) [arXiv:hep-th/0501098]; Phys. Rev. **D70**, 083514 (2004) [arXiv:hep-th/0406141].

[49] J. M. Cline, J. Descheneau, M. Giovannini and J. Vinet, JHEP **0306**, 048 (2003) [arXiv:hep-th/0304147].

[50] W. Israel, Nuovo Cim. **B44**, 1 (1966); Erratum-ibid. **B48**, 463 (1967).

[51] W. Israel, Bull, Am. Phys. Soc. **20**, 98 (1975); Phys. Rev. **D15**, 935 (1977).

[52] R. Geroch and J. Traschen, Phys. Rev. **D36**, 1017 (1987).

[53] V. P. Frolov, W. Israel and W. G. Unruh, Phys. Rev. **D39**, 1084 (1989).

[54] Work in progress. Preliminary consideration suggests stability at least near $\eta_\pm = 1$ at low energy, but detailed analysis remains to be done.

[55] C. Lanczos, Phys. Zeits, **23**, 539 (1922); Ann. der Phys. **74**, 518 (1924).

[56] T. Shiromizu, Y. Himemoto and K. Takahashi, Phys. Rev. **D70**, 107303 (2004) [hep-th/0405071]; T. Shiromizu, K. Takahashi, Y. Himemoto and S. Yamamoto, Phys. Rev. **D70**, 123524 (2004) [hep-th/0407268].

[57] K. Koyama and K. Koyama, "Gravitational backreaction of anti-D branes in the warped compactification", hep-th/0505256.